

# Signature-based Searches for New Physics in Diphoton Events; .... $[\gamma\gamma+X]$ ....

Sung-Won Lee & Ray Culbertson

#### Outline

Motivation

- Search Strategy
- First look in ~400 pb-1 diphoton data Diphoton Event Selection and Photon ID
- Searches for Anomalous Events and Results In this talk, I will focus on diphoton+X production, where X is MET, Jets
- Next to do...

## Motivation [I]


- The Standard Model has been enormously successful, but it is incomplete.
- Many theories/models attempt to address these issues, but none have been experimentally verified yet...
- Many reasons to believe there are new fundamental particles out there to be discovered at TeV scale.
- How does one search for new particles at the Tevatron?
  - → We know what Standard Model events look like.
  - → Look for events which are Un-Standard Model Like

# Motivation [II]

- Analysis goal is to find incontrovertible evidence for new physics beyond the Standard Model using as a photon(s) signature.
- There are a large number of important and well-motivated theoretical models which make a strong case for looking for new physics in events with a photon in the final state.
- These theories include SUSY, Extra Dimensions (ED), Grand Unified Theories, Composite models, Anomalous couplings and Higgs models.
  - SUSY  $(N_2 \rightarrow \gamma N_1, Light Gravitinos)$   $\longrightarrow$  mSUGRA
  - Extra Dimensions ADD & RS
  - Composite Models (Excited Leptons)
  - Technicolour: Photon+Dijet
  - Bosephilic Higgs: W/Z+Higgs  $\rightarrow$  W/Z+ $\gamma\gamma$
  - Anomalous  $W(\rightarrow \ell \nu)/Z(\rightarrow \ell \ell, \nu \nu) + \gamma$  production


**GMSB** 

# Example Signature - SUSY




# Search Strategy [I]

- Model-independent searches for "New Physics" in the photon final state.
- The basic signatures: Photon(s)+X(additional object(s)) in the final-state.



# Search Strategy [II]

- Model-independent searches for "New Physics" in the photon final state.
- The basic signatures: Photon(s)+X(additional object(s)) in the final-state.



## **Event Selection**

- New diphoton dataset [cdip0da\_diphoton]
  - equivalent to 5.3.3\_nt, dev\_242\_12 official prescriptions for calibrations, tagging, Zvertexing etc. [New Jet Correction]
- Global Event Selections
  - Two diphoton triggers in Central
  - $\bullet$   $|Z_{\text{vertex}}| < 60 \text{ cm}$
  - GoodRuns for diphoton analyses in 5.3.3 (435 pb<sup>-1</sup>)
- Standard Photon Identification and Isolation.
- Remove cosmic rays which brem or shower.
- Remove events caused by beam-related sources. [need to be updated; see highest MET event]
- Additional cuts to remove MET mis-measurement.

## Photon Identification

Baseline Analysis Cuts for High Pt Photons V2.1

<u>Variable</u>

• Had/EM

• Chi2 (Strips+Wires)/2.0

• N track (N3D)

Track Pt

• Cone 0.4 IsoEtCorr

• Cone 0.4 Track Iso

Fiducial

Cut

< 0.125 | | < 0.055 + 0.00045 \* ECorr

< 20

<=1

< 1+0.005\*EtCorr

EtCorr<20: < 0.1\*EtCorr

EtCorr>20: < 2.0+0.02\*(EtCorr-20.0)

< 2.0+0.005\*EtCorr

2nd CES cluster E\*sin(theta) (strip and wire)


EtCorr<18: < 0.14\*EtCorr

EtCorr>18: < 2.4+0.01\*EtCorr

Ces |X|<21 cm, 9< Ces |Z|<230 cm


Two isolated central photons with Et > 13 GeV 7295 diphoton events pass all candidate selesction requirement.

# [1] $\gamma\gamma+X$ Search, where X=MET



- Natural extension of previous analysis
   [CDF hep-ex/0410053]
- $4.11.2 \rightarrow 5.3.3$ \_nt
- 202 pb-1  $\rightarrow$  435 pb<sup>-1</sup>
- 5(3) events observed with MET>30(40) GeV
- trying to understand MET in 5.3.3\_nt
- revisit to Gaussian MET model using 5.3.3 Z sample
- optimize MET cut
- trying to reduce any noncollision backgrounds


#### Run 167998, Event 1041462






Et=(19,16), Met=39.9

#### Run 183617, Event 5889540




Et=(23,19), Met=36.2

#### Run 184518, Event 2144126



Et=(27,16), Met=50.8


#### Run 184868, Event 2536726





Et=(31,14), Met=54.2


#### Run 185009, Event 8965954



15

## [2] $\gamma\gamma$ +X Search, where X=Jets

- In some models
   Higgs couples only
   to vector bosons
- These are know as "Bosophilic" or "fermiophobic" Higgs
- CDF can search for γγ production in association with a W/Z boson.



 Look in γγ data for anomalous production on associated jets from quark decays of W's

## Next To Do...

- just start to look in ~400 pb<sup>-1</sup> diphoton data for new physics search using model-independent manner.
- Object Identifications
   eg. Jets, b-tags, c-tags,
   electrons, muons, and taus
- fake rate studies.
- Understanding of the signal and background systematically.

HOPE TO OPEN THE (SOME) BOX SOON ©

${ m E_T^{\gamma}} > 12~{ m GeV}~{ m Threshold}$		
Signature (Object)	Obs.	Expected
$E_{\rm T} > 35 \; { m GeV}, \;  \Delta \phi_{\rm r_{\rm r}-jet}  > 10^{\circ}$	Ţ	$0.5 \pm 0.1$
$ m N_{jet} \ge 4,  E_T^{jet} > 10  GeV,   \eta^{jet}  < 2.0$	2	$1.6 \pm 0.4$
Central e or $\mu$ , $\mathrm{E_T^{e \text{ or } \mu}} > 25 \text{ GeV}$	3	$0.3 \pm 0.1$
Central $\tau$ , $E_T^{\tau} > 25 \text{ GeV}$	1	$0.2 \pm 0.1$
$b$ -tag, $\mathrm{E}_{\mathrm{T}}^{b} > 25~\mathrm{GeV}$	2	$1.3 \pm 0.7$
Central $\gamma$ , $\mathrm{E_{T}^{\gamma_{3}}} > 25~\mathrm{GeV}$	0	$0.1 \pm 0.1$
${ m E_T^{\gamma} > 25~GeV~Threshold}$		
Object	Obs.	Exp.
$\mathbb{E}_{\mathrm{T}} > 25 \; \mathrm{GeV}, \;  \Delta \phi_{\mathbf{r}_{\mathrm{r}}-\mathrm{jet}}  > 10^{\circ}$	2	$0.5 \pm 0.1$
$ m N_{Jet} \ge 3, \ E_{T}^{Jet} > 10 \ GeV, \  \eta^{Jet}  < 2.0$	0	$1.7 \pm 1.5$
Central e or $\mu$ , $E_T^{e \text{ or } \mu} > 25 \text{ GeV}$	1	$0.1 \pm 0.1$
Central $\tau$ , $E_{\rm T}^{\tau} > 25 \; {\rm GeV}$	0	$0.03 \pm 0.0$
• / ±	0 0	$0.03 \pm 0.08$ $0.1 \pm 0.1$
Central $\tau$ , $E_{\rm T}^{\tau} > 25~{ m GeV}$	0 0 0	

Number of observed and expected  $\gamma\gamma$  events with additional objects in 85 pb<sup>-1</sup>